

Pollination Insights - Improving Pollination in Blueberry




Lisa W. DeVetter
Professor of Horticulture
Washington State University







Blueberry Pollination







- Most cultivars require or benefit from insect-mediated pollination
- Honey bees are the primary pollinator
- Pollination often variable and insufficient
- Growers need evidence-based strategies to improve blueberry pollination



Topics Highlighted Today



Pollination requirements



Honey bee pollination




Bumble bee pollination



Mitigating extreme heat




New technology (just a few words)





Pollination Requirements

- Highbush blueberry has a high degree of self-compatibility
- Yet, *Vaccinium* is partially self-sterile with an early-acting inbreeding depression
- Self-pollination causes some embryo abortion, which can limit fruit set and berry development
- Cross-pollination improves pollination outcomes across most blueberry cultivars





Cultivar Responses to Cross Pollination

Cultivar	Crossed with	Fruit mass	Firmness	Sugar:acid ratio	Ripening time
Arcadia	Arcadia Emerald Farthing	+	+	II	Up to 1 week faster
Colossus	Arcadia Emerald Farthing	+	+	=	Over 2 weeks faster
Farthing	Arcadia Emerald Farthing	+	=	=	Up to 8 days faster
Sentinel	Arcadia Emerald Farthing	+	+	=	Up to 9 days faster
Emerald	Arcadia Emerald Farthing	+	+	+	Up to a week faster
Optimus	Arcadia Emerald Farthing	+	+	+	Over 2 weeks faster


Data from Chabert and Mallinger at the University of Florida. Data based on greenhouse experiments and may not be transferrable to the field.




Breeding for Parthenocarpy









- Parthenocarpy = fruit development without fertilization (and pollination)
- Non-pollinated parthenocarpic fruits were smaller, sweeter, less firm, and took longer to develop but still met market standards
- Breeding for parthenocarpy possible

Genotypes by Treatment

Legend: Open Netted




Honey Bee Pollination

- Individual blueberry flower requires
 5-15 visits by a honey bee
- 25-274 pollen tetrads per flower to maximize fruit set and mass
- Low attraction to blueberry flowers, but they can get the job done!
- Hive placement and colony strength impacts foraging and pollination outcomes






Hive Placement Matters









Clumped



Dispersed

- Clumping hives led to a 2.6- and 1.3-fold increase in honey bee visitation in the Midwest and PNW, respectively, with no effect of stocking density
- No effects on fruit set, berry weight, nor seed set





Colony Strength (Worker Population Size) Varies











15,000 workers


30,000 workers

60,000 workers



It's Not Just Density – Colony Strength Counts!






- Increasing honey bee visitation improves pollination and yield outcomes (to a point)
- Visitation is best predicted by stocking strength (not field-level stocking density), underscoring the importance of stocking fields with strong colonies

Returning forager count



Stocking strength





- Honey bees are highly dependent on the sun for navigation
- Polytunnels reduce foraging and deplete hives when they fail to return to their colonies
- Middle sections of tunnels experience greater pollination deficits

Promoting Pollination in Protected Culture




- Stock with strong honey bee colonies (e.g., 6-10 hives/ha)
- Avoid UV-absorbing plastics and long tunnels (≥100 m)
- Place hives near tunnel ends
- Lift tunnel edges and leave ends open during bloom
- Possibly place hives in tunnels → rotate out with new hives every 2 weeks to maintain strong foraging populations
- Supplement honey bee nutrition
- Use additional pollinators (e.g., bumble bees)






Bumble Bee Pollination

- Buff-tailed bumble bee (*Bombus terrestris*)
- Good option for protected culture → less dependent on sunlight for navigation
- "Buzz pollination"
- More effective pollinator compared to honey bees (1 flower visit can lead to full pollination)
- Can be combined with honey bees
- Several commercial suppliers (Biobest, Koppert, Agrobio, etc.)






Optimizing Bumble Bee Pollination




= Pollination check points



- Order hives 8 weeks in advance
- Hives should arrive at ~5-10% bloom
- Single colony lasts 6-8 weeks before shifting into queen production
- Optimal hive density depends on colony size, flower density, and whether other pollinators are being used:
 - Start conservatively with a low density (9-10 hives/ha) and increase over time
 - Watch for over pollination
 - Work with your supplier

Images: Microsoft Copilot (2025)



- **Bumble Bee Deployment**
 - Deploy the day of arrival
 - Uncap sugar water (e.g., Biogluc®)
 - Do not place in direct light
 - Keep in a ventilated area (below 80% RH)
 - Place near pathways and distribute
 - Entrance/exit should be visible
 - Elevate 30-60 cm off the ground
 - Do not cluster more than 2 hives together
 - Protect from moisture
 - Use petroleum jelly if ants become problematic
 - Open hive after 15 minutes and monitor regularly
 - Dispose of hives after pollination is complete


Image: Microsoft Copilot (2025)



Extreme Heat Disrupts Pollination



- In vitro studies show pollen performance declines and is nonrecoverable as temperatures increase from 30-40°C
- Extreme heat events reduce pollen viability and functioning and can have nutritional effects that impact bee survival



Mean pollen tube length of four cultivars after 4 and 24 h of exposure to 10, 20, 30, or 40 °C.




Mitigating Extreme Heat









- Overhead cooling using micro-sprinklers in open fields reduced flower temperatures by 4 °C
- Honey bee activity reduced in 'Draper' only (yield data being analyzed)
- Current recommendation is to cool flowers when air temperatures are ≥ 29
 °C and to automate cooling cycles
- Also investigating biostimulants derived from kelp (Ascophyllum nodosum) and glycine betaine
- Research ongoing



Pollination Technology










- Pollination tech field is buzzing, but what is worth the investment?
- Partnership with universities –
 objective data to inform decisions
- Encouraging results with Beeflow
- Ongoing trials with BloomX





Pollination Resources











Blossoming Success:

BEST PRACTICES FOR NORTHERN HIGHBUSH BLUEBERRY POLLINATION

Salena Heimreich', Andony Melathopoulos', Zoe Little', Stan Chabert', Meghan O. Milbrath', Rachel E. Mallinger', Jenna Walters', Lauren Goldstein', Rutis Isaacs', Suzette P. Galinato', Clark Kogan', Emma Rogers', Kayla Brouwer', Maxime Erenets'', David Crowder', Lisa W. DeVetter'

Department of Entomology, Washington State University, Northwestern Washington Research and Extension

Center, Mount Vernon, WA, USA

University, Corvallis, OR, USA

3 Department of Entomology and Nematology, University of Florido Galnesville, FL, USA

4 Department of Entomology, Michigan State University, East Lansina, MI, US

5 Agriculture and Natural Resource Washington State University, Pull WA, USA

6 StatsCraft LLC Spokane WA US

7 Department of Horticulture, Was State University, Northwestern Washington Research and Exter Center, Mount Vernon, WA, USA

8 Department of Entomology, Washing Strete Linksprote, Bullman, W& LISA

Objective

This document describes on orthern highbus has blueberny (Voccinium corymbosum, the foreign refered to as "blueberny") politication and factors that promote politication of factors that promote politication of politications of the politication of the politication of the politication of polytiumete, etc., leg repensives, polytiumete, etc., leg repensives, polytiumete, etc., leg repensives polytiumete, etc., and politication of politication of the politication of politic

Defining pollination

Pollination is the movement of pollen between the reproductive parts of a flowering plant, which results in tertilization, production of seeds, and multi-development. In pollination-dependent crops, this pollen movement is facilitated by insacts, mainly bees. In bluebern, insact-vitalitation to flowers increase furth set, between plant set, between the production of the production of the production of the plant plant

























Conclusions

- Honey bees and bumble bees can be combined to improve pollination in protected culture, but further research is needed for optimization
- Important to establish good relationships with your bee supplier(s)
- Cross-pollination is beneficial for most cultivars
- Overhead cooling can reduce the negative effects of extreme heat
- Pollination tech is "buzzing," but require validation
- Morocco has >930 bee species → future research should explore how to safely integrate wild pollinators for more resilient pollination and enhance conservation efforts







Thank you! Questions?



lisa.devetter@wsu.edu





This work is supported by Specialty Crops Research Initiative Award #2020-02627 from the USDA National Institute of Food and Agriculture. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.













